antibodies .- online.com





Datasheet for ABIN873167 anti-Importin 9 antibody (AA 601-729)

| Coto | Product | nage |
|-------|---------|--------|
| 00 10 | Toduci | . payt |

| Overview | |
|-----------------------|--|
| Quantity: | 100 μL |
| Target: | Importin 9 (IPO9) |
| Binding Specificity: | AA 601-729 |
| Reactivity: | Mouse |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This Importin 9 antibody is un-conjugated |
| Application: | Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)) |
| Product Details | |
| Immunogen: | KLH conjugated synthetic peptide derived from human Importin 9 |
| Isotype: | IgG |
| Cross-Reactivity: | Mouse |
| Predicted Reactivity: | Human,Rat,Dog,Cow,Sheep,Pig,Horse,Chicken,Rabbit |
| Purification: | Purified by Protein A. |
| Target Details | |

Importin 9 (IPO9)

Target:

Target Details

| Alternative Name: | Importin 9 (IPO9 Products) | |
|---------------------|--|--|
| Background: | Synonyms: B cell antigen receptor Ig beta associated protein 1, BPM 90, BPM L, BPM-L, BPM90 | |
| | BPML, IBAP 1, Imp 9, Importin 9, Importin9, Importin-9, Novel centrosomal protein RanBPM, | |
| | RAN binding protein 9, Ran binding protein centrosomal, Ran binding protein M, Ran BP9, Ran- | |
| | binding protein 9, Ran-binding protein M, RANB9_HUMAN, RanBP 7, RANBP 9, RanBP7, RanBP9 RanBPM. | |
| | Background: May act as an adapter protein to couple membrane receptors to intracellular | |
| | signaling pathways. May be involved in signaling of ITGB2/LFA-1 and other integrins. Enhances | |
| | HGF-MET signaling by recruiting Sos and activating the Ras pathway. Involved in activation of | |
| | androgen and glucocorticoid receptor in the presence of their cognate hormones. Stabilizes | |
| | TP73 isoform Alpha, probably by inhibiting its ubiquitination, and increases its proapoptotic | |
| | activity. Inhibits the kinase activity of DYRK1A and DYRK1B. Inhibits FMR1 binding to | |
| | RNA. Tissue specificity: Ubiquitously expressed, with highest levels in testes, placenta, heart, | |
| | and muscle, and lowest levels in lung. Within the brain, expressed predominantly by neurons in | |
| | the gray matter of cortex, the granular layer of cerebellum and the Purkinje cells. | |
| Gene ID: | 10048 | |
| Pathways: | Protein targeting to Nucleus | |
| Application Details | | |
| Application Notes: | WB 1:300-5000 | |
| | ELISA 1:500-1000 | |
| | IHC-P 1:200-400 | |
| | IHC-F 1:100-500 | |
| | IF(IHC-P) 1:50-200 | |
| | IF(IHC-F) 1:50-200 | |
| | IF(ICC) 1:50-200 | |
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Concentration: | 1 μg/μL | |
| Buffer: | 0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol. | |
| | | |

Handling

| Preservative: | ProClin |
|--------------------|--|
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | 4 °C,-20 °C |
| Storage Comment: | Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. |
| Expiry Date: | 12 months |