# antibodies -online.com





# anti-PLEKHG6 antibody (AA 451-550)



Go to Product page

$\sim$					
	1/6	⊃r	1//	$\triangle$	٨/

Quantity:	100 μL
Target:	PLEKHG6
Binding Specificity:	AA 451-550
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PLEKHG6 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human MyoGEF
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Horse,Rabbit
Purification:	Purified by Protein A.

## **Target Details**

Target:	PLEKHG6
Alternative Name:	MyoGEF (PLEKHG6 Products)

### **Target Details**

Expiry Date:

rarget Details		
Background:	Synonyms: Myosin interacting guanine nucleotide exchange factor, PH domain containing family G member 6, Pleckstrin homology domain containing family G with RhoGef domain member 6, Pleckstrin homology domain-containing family G member 6, PLEKHG6_PKHG6_HUMAN  Background: MyoGEF functions as a guanine nucleotide exchange factor activating the small GTPase RHOA, which, in turn, induces myosin filament formation. It activates RHOG but not activate RAC1, or to a much lower extent than RHOA and RHOG. Part of a functional unit, involving MyoGEF, MYH10 and RHOA, at the cleavage furrow to advance furrow ingression during cytokinesis. In epithelial cells, required for the formation of microvilli and membrane ruffles on the apical pole. Along with EZR, required for normal macropinocytosis.	
Gene ID:	55200	
Application Details		
Application Notes:  Restrictions:	WB 1:300-5000  ELISA 1:500-1000  IHC-P 1:200-400  IHC-F 1:100-500  IF(IHC-P) 1:50-200  IF(ICC) 1:50-200  For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.	

12 months