

Datasheet for ABIN873239  
**anti-STRIP1 antibody (AA 255-360)**[Go to Product page](#)

## 1 Image

## Overview

Quantity:	100 µL
Target:	STRIP1
Binding Specificity:	AA 255-360
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This STRIP1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human FAM40A
Isotype:	IgG
Specificity:	There is a chance that this product will crossreact with PARD6G due to 75 % sequence similarity.
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Pig,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

## Target Details


Target:	STRIP1
Alternative Name:	Fam40a ( <a href="#">STRIP1 Products</a> )
Background:	Synonyms: FA40A_HUMAN, fam40a, FLJ14743, KIAA1761, MGC148091, Protein FAM40A, RP4-773N10.1.  Background: Belongs to the FAM40 family. 4 isoforms of the human protein are produced by alternative splicing.
Gene ID:	85369

## Application Details

Application Notes:	WB 1:300-5000 ELISA 1:500-1000 IHC-P 1:200-400 IHC-F 1:100-500 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months



#### Immunohistochemistry

**Image 1.** Formalin-fixed and paraffin embedded human brain labeled with Anti FAM40A Polyclonal Antibody, Unconjugated (ABIN873239) at 1:200 followed by conjugation to the secondary antibody and DAB staining