

Datasheet for ABIN873829

anti-ATP-Binding Cassette, Sub-Family B (MDR/TAP), Member 1B (ABCB1B) (AA 1051-1280) antibody (Alexa Fluor 555)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	ATP-Binding Cassette, Sub-Family B (MDR/TAP), Member 1B (ABCB1B)
Binding Specificity:	AA 1051-1280
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	Alexa Fluor 555
Application:	Flow Cytometry (FACS), Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human MDR1
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

Target Details

Target:	ATP-Binding Cassette, Sub-Family B (MDR/TAP), Member 1B (ABCB1B)
Alternative Name:	MDR1 (ABCB1B Products)
Background:	Synonyms: P-Glycoprotein, Multi Drug Resistance Associated Protein, ABCB1, ATP-binding

Target Details

cassette sub-family B MDR/TAP, member 1, ABC20, CD243, CD243 antigen, CLCS, Multidrug resistance protein 1, P glycoprotein 1, p-GP, gp170, P gp, PGY1.

Background: P Glycoprotein, the product of the MDR1 gene, is expressed in distinct non-malignant cells, typically cells with secretory and excretory functions. It is assumed to function as an ATP-dependent drug efflux pump with broad substrate specificity. The highest expression of P Glycoprotein has been observed in kidney (proximal tubules), liver (bile canaliculi), adrenal gland and intestine, suggesting that the primary role of P Glycoprotein is in the normal secretion of physiological metabolites and ingested chemicals into bile, urine and the lumen of the intestinal tract. Elevated levels of P Glycoprotein have also been reported in multidrug-resistant cell lines and in colon, endometrial, ovarian, and breast tumors, as well as in sarcomas and leukemias / lymphomas.

Gene ID:	5243
----------	------

Application Details

Application Notes:	FCM 1:20-100 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
--------------------	--

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Liquid
---------	--------

Concentration:	1 µg/µL
----------------	---------

Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
---------	---

Preservative:	ProClin
---------------	---------

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
--------------------	--

Storage:	-20 °C
----------	--------

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
------------------	---

Expiry Date:	12 months
--------------	-----------