

## Datasheet for ABIN877160

## anti-FXYD1 antibody (pSer83) (AbBy Fluor® 350)



Go to Product page

Overview	
Quantity:	100 μL
Target:	FXYD1
Binding Specificity:	pSer83
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FXYD1 antibody is conjugated to AbBy Fluor® 350
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
Immunogen:	KLH conjugated synthetic phosphopeptide derived from human FXYD1 around the phosphorylation site of Ser83
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Pig
Purification:	Purified by Protein A.
Target Details	
Target:	FXYD1
Alternative Name:	FXYD1 (FXYD1 Products)

## **Target Details**

Background:	Synonyms: FXYD1 phospho S83, FXYD1 phospho Ser83, p-FXYD1 phospho S83, FXYD domain
	containing ion transport regulator 1, Phospholemman, PLM, fxyd1, PLM_HUMAN, FXYD
	domain-containing ion transport regulator 1.
	Background: PLM (FXYD1)is a member of a family of small membrane proteins that share a 35-
	amino acid signature sequence domain, beginning with the sequence PFXYD and containing 7
	invariant and 6 highly conserved amino acids. FXYD2, also known as the gamma subunit of the
	Na,K-ATPase, regulates the properties of that enzyme. FXYD1 (phospholemman), FXYD2
	(gamma), FXYD3 (MAT-8), FXYD4 (CHIF), and FXYD5 (RIC) have been shown to induce channel
	activity in experimental expression systems. PLM may be phosphorylated by several kinases,
	including protein kinase A, protein kinase C, NIMA kinase, and myotonic dystrophy kinase. It is
	thought to form an ion channel or regulate ion channel activity.
Gene ID:	5348
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months