

Datasheet for ABIN877835

anti-CDC27 antibody (pThr244) (AbBy Fluor® 488)



Overview

Overview	
Quantity:	100 μL
Target:	CDC27
Binding Specificity:	pThr244
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CDC27 antibody is conjugated to AbBy Fluor® 488
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded
	Sections) (IF (p))
Product Details	
Immunogen:	KLH conjugated synthetic phosphopeptide derived from human CDC27 around the
	phosphorylation site of Thr244
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human,Dog,Pig,Horse,Rabbit
Purification:	Purified by Protein A.
Target Details	
Target:	CDC27

Target Details

Alternative Name:	CDC27 (CDC27 Products)
Background:	Synonyms: ANAPC3, Anaphase Promoting Complex 3, Anaphase promoting complex protein 3,
	Anaphase Promoting Complex Subunit 3, Anaphase-promoting complex subunit 3, APC 3,
	APC3, APC-3, Cdc 27, Cdc27, Cdc-27, CDC27 homolog, CDC27_HUMAN, CDC27Hs, Cell division
	cycle 27, Cell division cycle protein 27 homolog, H NUC, H-NUC, HNUC, NUC2 homolog.
	Background: The protein encoded by this gene shares strong similarity with Saccharomyces
	cerevisiae protein Cdc27, and the gene product of Schizosaccharomyces pombe nuc 2. This
	protein is a component of anaphase-promoting complex (APC), which is composed of eight
	protein subunits and highly conserved in eucaryotic cells. APC catalyzes the formation of cyclin
	B-ubiquitin conjugate that is responsible for the ubiquitin-mediated proteolysis of B-type cyclins
	This protein and 3 other members of the APC complex contain the TPR (tetratricopeptide
	repeat), a protein domain important for protein-protein interaction. This protein was shown to
	interact with mitotic checkpoint proteins including Mad2, p55CDC and BUBR1, and thus may be
	involved in controlling the timing of mitosis. Alternatively spliced transcript variants encoding
	different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]
Gene ID:	996
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.

Handling

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months