## antibodies .- online.com





## anti-INPPL1 antibody (pTyr1135) (AbBy Fluor® 350)



$\sim$			
	$  \backslash / \cap$	r\/I	$I \cap V$

Overview		
Quantity:	100 μL	
Target:	INPPL1	
Binding Specificity:	pTyr1135	
Reactivity:	Human, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This INPPL1 antibody is conjugated to AbBy Fluor® 350	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	
Product Details		
Immunogen:	KLH conjugated synthetic phosphopeptide derived from human SHIP2 around the phosphorylation site of Tyr1135	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse	
Predicted Reactivity:	Dog,Cow,Horse,Rabbit	
Purification:	Purified by Protein A.	
Target Details		
Target:	INPPL1	

## **Target Details**

Alternative Name:	INPPL1 (INPPL1 Products)	
Background:	Synonyms: 4, 5-trisphosphate 5-phosphatase 2, 51C protein, EC 3.1.3.n1, inositol	
	polyphosphate phosphatase like 1, Inositol polyphosphate phosphatase like protein 1, Inositol	
	polyphosphate phosphatase-like protein 1, INPPL-1, INPPL1, Phosphatidylinositol 3,	
	Phosphatidylinositol 3,4,5 trisphosphate 5 phosphatase 2, Protein 51C, SH2 domain containing	
	inositol 5' phosphatase 2, SH2 domain-containing inositol 5"-phosphatase 2, SH2 domain-	
	containing inositol phosphatase 2, SHIP-2, SHIP2, SHIP2_HUMAN.	
	Background: The steady state of protein tyrosyl phosphorylation in cells is regulated by the	
	opposing action of tyrosine kinases and protein tyrosine phosphatases (PTPs). Several groups	
	have independently identified a non transmembrane PTP, designated SHPTP1 (also known as	
	PTP1C, HCP and SHP), which is primarily expressed in hematopoietic cells and characterized	
	by the presence of two SH2 domains N terminal to the PTP domain. A second and much more	
	widely expressed PTP with SH2 domains, SHPTP2 (also designated PTP1D and Syp), has been	
	identified. SHP2 is a protein tyrosine phosphatase that is widely expressed and plays a	
	regulatory role in various cell signaling events that are important for many cell functions, such	
	as mitogenic activation, metabolic control, transcription regulation, and cell migration.	
Gene ID:	3636	
Pathways:	Platelet-derived growth Factor Receptor Signaling	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	

## Handling

	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months