antibodies .- online.com







anti-DLG2 antibody (pTyr340) (Alexa Fluor 555)



()	11/0	K\ /	iew	1
	\cup	'I V/I	$\square \vee \vee$	ı

Overview	
Quantity:	100 μL
Target:	DLG2
Binding Specificity:	pTyr340
Reactivity:	Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DLG2 antibody is conjugated to Alexa Fluor 555
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
lmmunogen:	KLH conjugated synthetic phosphopeptide derived from human PSD93 around the phosphorylation site of Tyr340
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Dog,Horse,Chicken
Purification:	Purified by Protein A.
Target Details	
Target:	DLG2

Target Details

Alternative Name:	PSD93 (DLG2 Products)	
Background:	Synonyms: Channel associated protein of synapse 110, Channel associated protein of	
	synapses 110kD, Channel-associated protein of synapse-110, Chapsyn 110, Chapsyn-110,	
	Chapsyn110, discs large homolog 2, Discs, large homolog 2 Drosophila, Disks large homolog 2,	
	DKFZp781D1854, DKFZp781E0954, Dlg 2, dlg2, DLG2_HUMAN, Dlgh 2, Dlgh2, FLJ37266,	
	Gm1197, MGC131811, Postsynaptic density protein PSD 93, Postsynaptic density protein PSD-	
	93, Postsynaptic density protein PSD93, PSD 93, PSD93.	
	Background: PSD 93 is believed to participate in the clustering of certain proteins, including N-	
	methyl-D-aspartate (NMDA) receptors and shaker-type potassium channels at the synaptic	
	membrane. There are two principal modes of interaction between PSD 93 and other proteins.	
	NMDA receptors and shaker-type potassium channels both share C-terminal sequence	
	homology consisting of a threonine/serine-X-valine-COOH (T/SXV) motif. Other neuronal	
	proteins that share this motif (beta 1 adrenergic receptor, some serotonin receptors, some	
	sodium channel subunits, and additional potassium channel subunits) may interact with PSD	
	93 by binding to its PDZ domains. Neuronal nitric oxide synthase (nNOS), which lacks the	
	T/SXV motif but which has its own PDZ domain, has been shown to associate with PSD 93 in	
	vitro through a pseudo-homotypic PDZ-PDZ interaction.	
Gene ID:	1740	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
	Liquid	
Handling Format: Concentration:	Liquid 1 μg/μL	
Format: Concentration:	<u> </u>	
Format:	1 μg/μL	
Format: Concentration:	1 μg/μL Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	

Handling

	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months