



[Go to Product page](#)

Datasheet for ABIN880435

anti-INPPL1 antibody (pTyr986, pTyr987) (AbBy Fluor® 350)

Overview

Quantity:	100 µL
Target:	INPPL1
Binding Specificity:	pTyr986, pTyr987
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This INPPL1 antibody is conjugated to AbBy Fluor® 350
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic phosphopeptide derived from human SHIP2 around the phosphorylation site of Tyr986/987
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Pig, Horse, Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	INPPL1
Alternative Name:	INPPL1 + (INPPL1 Products)

Target Details

Background: Synonyms: 4, 5-trisphosphate 5-phosphatase 2, 51C protein, EC 3.1.3.n1, inositol polyphosphate phosphatase like 1, Inositol polyphosphate phosphatase like protein 1, Inositol polyphosphate phosphatase-like protein 1, INPPL-1, INPPL1, Phosphatidylinositol 3, Phosphatidylinositol 3,4,5 trisphosphate 5 phosphatase 2, Protein 51C, SH2 domain containing inositol 5' phosphatase 2, SH2 domain-containing inositol 5"-phosphatase 2, SH2 domain-containing inositol phosphatase 2, SHIP-2, SHIP2, SHIP2_HUMAN.

Background: The steady state of protein tyrosyl phosphorylation in cells is regulated by the opposing action of tyrosine kinases and protein tyrosine phosphatases (PTPs). Several groups have independently identified a non transmembrane PTP, designated SHPTP1 (also known as PTP1C, HCP and SHP), which is primarily expressed in hematopoietic cells and characterized by the presence of two SH2 domains N terminal to the PTP domain. A second and much more widely expressed PTP with SH2 domains, SHPTP2 (also designated PTP1D and Syp), has been identified. SHP2 is a protein tyrosine phosphatase that is widely expressed and plays a regulatory role in various cell signaling events that are important for many cell functions, such as mitogenic activation, metabolic control, transcription regulation, and cell migration.

Gene ID: 3636

Pathways: [Platelet-derived growth Factor Receptor Signaling](#)

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months