antibodies .- online.com







anti-APEH antibody (AA 101-200) (Alexa Fluor 350)



| () | 1/0 | r\ / I | 014 | |
|-----|-----|--------|-----|---|
| () | ve | I V I | -v | V |
| | | | | |

| Quantity: | 100 μL |
|----------------------|--|
| Target: | APEH |
| Binding Specificity: | AA 101-200 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This APEH antibody is conjugated to Alexa Fluor 350 |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human AARE |
|-----------------------|--|
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Predicted Reactivity: | Mouse,Rat,Cow,Sheep,Pig,Horse,Rabbit |
| Purification: | Purified by Protein A. |

Target Details

| Target: | APEH |
|-------------------|----------------------|
| Alternative Name: | AARE (APEH Products) |

Target Details

| Background: | Synonyms: AARE, ACPH, Acyl peptide hydrolase, Acylamino acid releasing enzyme, | |
|---------------------|---|--|
| | Acylaminoacyl peptidase, APEH, APH, D3F15S2, D3S48E, DNF15S2, N acylaminoacyl peptide | |
| | hydrolase, OPH, Oxidized protein hydrolase, ACPH_HUMAN. | |
| | Background: AARE (Acylamino-acid-releasing enzyme) is also known as Acyl-peptide hydrolase. | |
| | It catalyzes the hydrolysis of the terminal acetylated amino acid preferentially from small | |
| | acetylated peptides. The acetyl amino acid formed by this hydrolase is further processed to | |
| | acetate and a free amino acid by an aminoacylase. It can play an important role in destroying | |
| | oxidatively damaged proteins in living cells. Deletions of this gene locus corresponding to the | |
| | protein are found in various types of carcinomas, including small cell lung carcinoma and renal | |
| | cell carcinoma. | |
| Gene ID: | 327 | |
| Application Details | | |
| Application Notes: | IF(IHC-P) 1:50-200 | |
| | IF(IHC-F) 1:50-200 | |
| | IF(ICC) 1:50-200 | |
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Concentration: | 1 μg/μL | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and | |
| | 50 % Glycerol. | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be | |
| | handled by trained staff only. | |
| Storage: | -20 °C | |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. | |
| Expiry Date: | 12 months | |
| . , | | |