





anti-Adenylosuccinate Lyase antibody (AA 185-280) (Biotin)



Go to Product page

()	۱ ۱	\cap	r	/1	\cap	۱ ۸	1
0	'V	ㄷ	I١	νı	\Box	V	۷

Quantity:	100 μL	
Target:	Adenylosuccinate Lyase (ADSL)	
Binding Specificity:	AA 185-280	
Reactivity:	Human, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Adenylosuccinate Lyase antibody is conjugated to Biotin	
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Adenylosuccinate Lyase	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse	
Predicted Reactivity:	Rat,Dog,Cow,Pig,Horse	
Purification:	Purified by Protein A.	

Target Details

Target:	Adenylosuccinate Lyase (ADSL)
Alternative Name:	Adenylosuccinate Lyase (ADSL Products)

Target Details	
Background:	Synonyms: Adenylosuccinase, Adenylosuccinate lyase, ADSL, AMPS, ASase, ASL, PUR8_HUMAN.
	Background: Adenylsuccinate lyase is involved in both de novo synthesis of purines and
	formation of adenosine monophosphate from inosine monophosphate. It catalyzes two
	reactions in AMP biosynthesis: the removal of a fumarate from succinylaminoimidazole
	carboxamide (SAICA) ribotide to give aminoimidazole carboxamide ribotide (AICA) and removal
	of fumarate from adenylosuccinate to give AMP. Adenylosuccinase deficiency results in
	succinylpurinemic autism, psychomotor retardation, and , in some cases, growth retardation
	associated with muscle wasting and epilepsy. Two transcript variants encoding different
	isoforms have been found for this gene. [provided by RefSeq, Jul 2008].
Pathways:	Ribonucleoside Biosynthetic Process
Application Details	
Application Notes:	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months