

## Datasheet for ABIN881696

Overview

## anti-ADRB1 antibody (AA 181-250) (AbBy Fluor® 488)



- Overview	
Quantity:	100 μL
Target:	ADRB1
Binding Specificity:	AA 181-250
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ADRB1 antibody is conjugated to AbBy Fluor® 488
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Flow Cytometry (FACS)
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human ADRB1

immunogen:	KLH conjugated synthetic peptide derived from numan ADRB i
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Pig
Purification:	Purified by Protein A.

## Target Details

Target:	ADRB1
Alternative Name:	ADRB1 (ADRB1 Products)

## **Target Details**

Background:	Synonyms: RHR, B1AR, ADRB1R, BETA1AR, Beta-1 adrenergic receptor, Beta-1 adrenoreceptor,
background.	Beta-1 adrenoceptor, ADRB1
	Background: Beta-adrenergic receptors mediate the catecholamine-induced activation of
	adenylate cyclase through the action of G proteins. This receptor binds epinephrine and
	norepinephrine with approximately equal affinity. Mediates Ras activation through G(s)-alpha-
	and cAMP-mediated signaling.
Gene ID:	153
UniProt:	P08588
Pathways:	cAMP Metabolic Process, Cellular Glucan Metabolic Process, Regulation of Muscle Cell
	Differentiation, Synaptic Membrane, Regulation of G-Protein Coupled Receptor Protein
	Signaling, G-protein mediated Events, Interaction of EGFR with phospholipase C-gamma, Brown
	Fat Cell Differentiation
	Fat Celi Differentiation
Application Details	
Application Notes:	FCM 1:20-100
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months