antibodies - online.com







anti-MTDH antibody (AA 131-230) (Alexa Fluor 488)



| () | 1 / | \sim | KI / | 110 | Νę |
|-----|-----|----------|------|-----|-------|
| | 1// | \vdash | I \/ | 1 ← | ٠// ٢ |
| | | | | | |

| Quantity: | 100 μL | |
|----------------------|---|--|
| Target: | MTDH | |
| Binding Specificity: | AA 131-230 | |
| Reactivity: | Human | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This MTDH antibody is conjugated to Alexa Fluor 488 | |
| Application: | Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) | |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human AEG-1 |
|-----------------------|---|
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Predicted Reactivity: | Mouse,Rat,Dog,Cow,Pig |
| Purification: | Purified by Protein A. |

Target Details

| Target: | MTDH |
|-------------------|-----------------------------|
| Alternative Name: | Aeg-1/Lyric (MTDH Products) |

Target Details

| Synonyms: 3D3, AEG1, AEG-1, LYRIC, LYRIC/3D3, Protein LYRIC, 3D3/LYRIC, Astrocyte elevated | |
|---|--|
| gene-1 protein, Lysine-rich CEACAM1 co-isolated protein, Metadherin, Metastasis adhesion | |
| protein, MTDH | |
| Background: Downregulates SLC1A2/EAAT2 promoter activity when expressed ectopically. | |
| Activates the nuclear factor kappa-B (NF-kappa-B) transcription factor. Promotes anchorage- | |
| independent growth of immortalized melanocytes and astrocytes which is a key component in | |
| tumor cell expansion. Promotes lung metastasis and also has an effect on bone and brain | |
| metastasis, possibly by enhancing the seeding of tumor cells to the target organ endothelium. | |
| Induces chemoresistance. | |
| 92140 | |
| Q86UE4 | |
| Cellular Response to Molecule of Bacterial Origin, Cell-Cell Junction Organization | |
| | |
| FCM 1:20-100 | |
| IF(IHC-P) 1:50-200 | |
| IF(IHC-F) 1:50-200 | |
| IF(ICC) 1:50-200 | |
| For Research Use only | |
| | |
| Liquid | |
| 1 μg/μL | |
| Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and | |
| 50 % Glycerol. | |
| ProClin | |
| This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be | |
| handled by trained staff only. | |
| -20 °C | |
| Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. | |
| 12 months | |
| | |