# antibodies -online.com





# anti-ANKRD11 antibody (AA 731-820) (Alexa Fluor 488)



Go to Product page

$\sim$			
	N/P	r\/	i⊢₩

Quantity:	100 μL	
Target:	ANKRD11	
Binding Specificity:	AA 731-820	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This ANKRD11 antibody is conjugated to Alexa Fluor 488	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human ANKRD11
Isotype:	lgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Horse
Purification:	Purified by Protein A.

## **Target Details**

Target:	ANKRD11
Alternative Name:	ANKRD11 (ANKRD11 Products)

## **Target Details**

Expiry Date:

rarget Details		
Background:	Synonyms: ANCO 1, ANCO1, Ankyrin repeat containing cofactor 1, Ankyrin repeat domain 11, Ankyrin repeat domain containing protein 11, LZ16, T13.  Background: Ankyrin is a membrane protein that mediates the attachment of the erythrocyte membrane skeleton to the plasma membrane and interacts with CD44 and inositol triphosphate. It contains three functional domains: a conserved N-terminal ankyrin repeat	
	domain (ARD(consisting of 2224 tandem repeats of 33 amino acids), a spectrin binding domain and a variably sized C-terminal regulatory domain. The ankyrin repeat is a 33-residue motif in	
	proteins consisting of two alpha helices separated by loops. It has been studied using multiple	
	sequence alignment to determine which conserved amino acid residues are critical for folding	
	and stability. Ankyrin-repeat proteins have been associated with a number of human diseases,	
	most notably, the cell cycle inhibitor p16 is associated with cancer and the Notch protein is a	
	key component of cell signaling pathways whose intracellular repeat domain is disrupted in	
	mutations that give rise to the neurological disorder known as CADASIL.	
Gene ID:	29123	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	

12 months