antibodies .- online.com





Datasheet for ABIN883766

anti-ASB3 antibody (AA 21-120) (HRP)

Overview 100 μL Quantity: ASB3 Target: Binding Specificity: AA 21-120 Reactivity: Human, Mouse Host: Rabbit Clonality: Polyclonal This ASB3 antibody is conjugated to HRP Conjugate: Application: ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)) **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human ASB3
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat,Dog,Cow,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	ASB3
Alternative Name:	ASB3 (ASB3 Products)

Target Details

Background:

Synonyms: Ankyrin repeat and SOCS box containing 3, Ankyrin repeat and SOCS box containing protein 3, Ankyrin repeat and SOCS box protein 3, ASB-3, ASB3_HUMAN, Weakly similar to ankyrin brain variant 2.

Background: Members of the suppressor of cytokine signaling (SOCS) family of proteins contain C-terminal regions of homology called the SOCS box, which serves to couple SOCS proteins and their binding partners with the elongin B and C complex. Serveral other families of proteins also contain SOCS boxes but differ from the SOCS proteins in the type of domain they contained upstream of the SOCS box. Four members of the ankyrin repeat and SOCS boxcontaining (ASB) protein family are identified and termed as ASB-1, ASB-2, ASB-3 and ASB-4. ASB-1 is expressed in multiple organs, including the hematopoietic compartment. ASB-1 knockout mice display a diminution of spermatogenesis with less complete filling of seminiferous tubules. Asb-2 is a novel retinoic-acid (RA)-induced gene in acute promyelocytic leukemia (APL) cells and its expression induces growth-inhibition and chromatin condensation recapitulating early events critical to RA-induced differentiaiton of APL cells. ASB-2 is directly induced by all-trans retinotic acid, by the binding of RARa to the RAR binding element/RXR binding element in the Asb-2 promoter.

Gene ID:

51130

Application Details

Δn	plication	Notes:
\neg	piicatioi	i NOLCS.

IHC-P 1:200-400

IHC-F 1:100-500

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish

Handling

	peroxidase.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months