



[Go to Product page](#)

Datasheet for ABIN883772

anti-ASB3 antibody (AA 21-120) (Alexa Fluor 647)

Overview

| | |
|----------------------|---|
| Quantity: | 100 µL |
| Target: | ASB3 |
| Binding Specificity: | AA 21-120 |
| Reactivity: | Human, Mouse |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This ASB3 antibody is conjugated to Alexa Fluor 647 |
| Application: | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| | |
|-----------------------|--|
| Immunogen: | KLH conjugated synthetic peptide derived from human ASB3 |
| Isotype: | IgG |
| Cross-Reactivity: | Human, Mouse |
| Predicted Reactivity: | Rat,Dog,Cow,Rabbit |
| Purification: | Purified by Protein A. |

Target Details

| | |
|-------------------|--|
| Target: | ASB3 |
| Alternative Name: | ASB3 (ASB3 Products) |

Target Details

| | |
|-------------|--|
| Background: | <p>Synonyms: Ankyrin repeat and SOCS box containing 3, Ankyrin repeat and SOCS box containing protein 3, Ankyrin repeat and SOCS box protein 3, ASB-3, ASB3, ASB3_HUMAN, Weakly similar to ankyrin brain variant 2.</p> <p>Background: Members of the suppressor of cytokine signaling (SOCS) family of proteins contain C-terminal regions of homology called the SOCS box, which serves to couple SOCS proteins and their binding partners with the elongin B and C complex. Several other families of proteins also contain SOCS boxes but differ from the SOCS proteins in the type of domain they contained upstream of the SOCS box. Four members of the ankyrin repeat and SOCS box-containing (ASB) protein family are identified and termed as ASB-1, ASB-2, ASB-3 and ASB-4. ASB-1 is expressed in multiple organs, including the hematopoietic compartment. ASB-1 knock-out mice display a diminution of spermatogenesis with less complete filling of seminiferous tubules. Asb-2 is a novel retinoic-acid (RA)-induced gene in acute promyelocytic leukemia (APL) cells and its expression induces growth-inhibition and chromatin condensation recapitulating early events critical to RA-induced differentiation of APL cells. ASB-2 is directly induced by all-trans retinoic acid, by the binding of RARα to the RAR binding element/RXR binding element in the Asb-2 promoter.</p> |
|-------------|--|

| | |
|----------|-------|
| Gene ID: | 51130 |
|----------|-------|

Application Details

| | |
|--------------------|--|
| Application Notes: | IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 |
|--------------------|--|

| | |
|---------------|-----------------------|
| Restrictions: | For Research Use only |
|---------------|-----------------------|

Handling

| | |
|--------------------|--|
| Format: | Liquid |
| Concentration: | 1 µg/µL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |

Handling

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months