

Datasheet for ABIN884409

anti-BAFF antibody (AA 221-309) (AbBy Fluor® 350)



Overview

| Overview | |
|-----------------------|--|
| Quantity: | 100 μL |
| Target: | BAFF (TNFSF13B) |
| Binding Specificity: | AA 221-309 |
| Reactivity: | Human, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This BAFF antibody is conjugated to AbBy Fluor® 350 |
| Application: | Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |
| Product Details | |
| Immunogen: | KLH conjugated synthetic peptide derived from mouse BAFF |
| Isotype: | IgG |
| Cross-Reactivity: | Human, Rat |
| Predicted Reactivity: | Mouse,Cow,Pig,Rabbit |
| Purification: | Purified by Protein A. |
| Target Details | |
| Target: | BAFF (TNFSF13B) |
| Alternative Name: | Baff (TNFSF13B Products) |

Target Details

| Background: | Synonyms: ApoL related ligand TALL 1, B cell Activating Factor, B lymphocyte stimulator , BLYS, |
|---------------------|---|
| | CD 257, CD257, CD257 antigen, delta BAFF, Dendritic cell derived TNF like molecule, DTL, DTL |
| | precursor, TALL 1, TALL1, THANK, TNF and APOL related leukocyte expressed ligand 1, TNF |
| | homolog that activates apoptosis, TNF homolog that activates apoptosis NKFB and JNK, |
| | TNFSF13B, TNFSF20, Tumor necrosis factor ligand superfamily member 13B, Tumor necrosis |
| | factor ligand superfamily member 20, Tumor necrosis factor like protein ZTNF4, Tumor |
| | necrosis factor superfamily member 13B, ZTNF4, TN13B_MOUSE. |
| | Background: The protein encoded by this gene is a cytokine that belongs to the tumor necrosis |
| | factor (TNF) ligand family. This cytokine is a ligand for receptors TNFRSF13B/TACI, |
| | TNFRSF17/BCMA, and TNFRSF13C/BAFFR. This cytokine is expressed in B cell lineage cells, |
| | and acts as a potent B cell activator. It has been also shown to play an important role in the |
| | proliferation and differentiation of B cells. [provided by RefSeq]. |
| Gene ID: | 24099 |
| UniProt: | Q9WU72 |
| Pathways: | NF-kappaB Signaling, Production of Molecular Mediator of Immune Response |
| Application Details | |
| Application Notes: | FCM 1:20-100 |
| | IF(IHC-P) 1:50-200 |
| | IF(IHC-F) 1:50-200 |
| | IF(ICC) 1:50-200 |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and |
| | 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be |
| | handled by trained staff only. |
| Storage: | -20 °C |
| | |

Handling

| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
|------------------|---|
| Expiry Date: | 12 months |