antibodies - online.com





Datasheet for ABIN884508

anti-BARD1 antibody (AA 101-200) (Alexa Fluor 555)



()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μL	
Target:	BARD1	
Binding Specificity:	AA 101-200	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This BARD1 antibody is conjugated to Alexa Fluor 555	
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human BARD1
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Pig,Horse,Chicken
Purification:	Purified by Protein A.

Target Details

Target:	BARD1
Alternative Name:	BARD1 (BARD1 Products)

Target Details

Synonyms: BRCA1-associated RING domain protein 1, BARD-1, BARD1	
Background: Probable E3 ubiquitin-protein ligase. The BRCA1-BARD1 heterodimer specifically	
mediates the formation of 'Lys-6'-linked polyubiquitin chains and coordinates a diverse range of	
cellular pathways such as DNA damage repair, ubiquitination and transcriptional regulation to	
maintain genomic stability. Plays a central role in the control of the cell cycle in response to	
DNA damage. Acts by mediating ubiquitin E3 ligase activity that is required for its tumor	
suppressor function. Also forms a heterodimer with CSTF1/CSTF-50 to modulate mRNA	
processing and RNAP II stability by inhibiting pre-mRNA 3' cleavage.	
580	
Q99728	
DNA Damage Repair	
FCM 1:20-100	
IF(IHC-P) 1:50-200	
IF(IHC-F) 1:50-200	
IF(ICC) 1:50-200	
For Research Use only	
Liquid	
1 μg/μL	
Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
50 % Glycerol.	
ProClin	
This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
handled by trained staff only.	
-20 °C	
Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
12 months	