antibodies -online.com





anti-BCCIP antibody (AA 75-180) (AbBy Fluor® 555)



Go to Product page

()	11/0	K\ /	iew
	\cup	'I V/I	$I \cap VV$

Quantity:	100 μL	
Target:	BCCIP	
Binding Specificity:	AA 75-180	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This BCCIP antibody is conjugated to AbBy Fluor® 555	
Application:	Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human BCCIP/TOK1
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Horse
Purification:	Purified by Protein A.

Target Details

Target:	BCCIP
Alternative Name:	BCCIP (BCCIP Products)
Background: Synonyms: bccip, BCCIP_HUMAN, BCCIPalpha, BCCIPbeta, BRCA2 and CDKN1A interactions and action of the control of th	

Preservative:

Storage:

Expiry Date:

Precaution of Use:

Storage Comment:

Target Details	
	protein, BRCA2 and CDKN1A-interacting protein, BRCA2 and Cip1/p21 interacting protein, Cdk inhibitor p21 binding protein, P21 and CDK associated protein 1, P21- and CDK-associated protein 1, Protein TOK-1, Protein TOK1, TOK 1, TOK 1alpha, TOK 1beta, TOK1. Background: FunctionMay promote cell cycle arrest by enhancing the inhibition of CDK2 activity by CDKN1A. May be required for repair of DNA damage by homologous recombination in conjunction with BRCA2. May not be involved in non-homologous end joining.
Gene ID:	56647
Application Details	
Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and

This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

50 % Glycerol.

handled by trained staff only.

ProClin

-20 °C

12 months