

[Go to Product page](#)

Datasheet for ABIN885426

anti-CTCFL antibody (AA 351-450) (Alexa Fluor 350)

Overview

Quantity:	100 µL
Target:	CTCFL
Binding Specificity:	AA 351-450
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CTCFL antibody is conjugated to Alexa Fluor 350
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human BORIS/CT27
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Pig, Horse
Purification:	Purified by Protein A.

Target Details

Target:	CTCFL
Alternative Name:	CT27 (CTCFL Products)
Background:	Synonyms: BORIS like protein, Brother of the regulator of imprinted sites, Cancer/testis antigen

Target Details

27, CCCTC binding factor zinc finger protein like, CCCTC-binding factor, CT27, CTCF paralog, CTCF T, CTCF-like protein, CTCFL, CTCFL_HUMAN, HMG 1L1, HMGB1L1, Putative high mobility group protein 1 like 1, Putative high mobility group protein B1 like 1, Transcriptional repressor CTCFL, Zinc finger protein CTCF-T.

Background: Testis-specific DNA binding protein responsible for insulator function, nuclear architecture and transcriptional control, which probably acts by recruiting epigenetic chromatin modifiers. Plays a key role in gene imprinting in male germline, by participating in the establishment of differential methylation at the IGF2/H19 imprinted control region (ICR). Directly binds the unmethylated H19 ICR and recruits the PRMT7 methyltransferase, leading to methylate histone H4 'Arg-3' to form H4R3sme2. This probably leads to recruit de novo DNA methyltransferases at these sites (By similarity). Seems to act as tumor suppressor. In association with DNMT1 and DNMT3B, involved in activation of BAG1 gene expression by binding to its promoter. Required for dimethylation of H3 lysine 4 (H3K4me2) of MYC and BRCA1 promoters.

Gene ID: 140690

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Handling

Expiry Date: 12 months