## antibodies -online.com





## anti-Protein Rta/BRLF1 antibody (AA 81-180) (Biotin)



| _ |   |   |    |    |   |   |   |
|---|---|---|----|----|---|---|---|
| 0 | V | е | r١ | /1 | е | V | I |

| Quantity:   | 100 μL   |  |
|---|--|--|
| Target:   | Protein Rta/BRLF1 (BRLF1)  |  |
| Binding Specificity:  | AA 81-180  |  |
| Reactivity:   | Virus, Human   |  |
| Host:   | Rabbit   |  |
| Clonality:  | Polyclonal   |  |
| Conjugate:  | This Protein Rta/BRLF1 antibody is conjugated to Biotin  |  |
| Application:  | Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), ELISA      |  |
| Product Details   |  |  |
| Immunogen:  | KLH conjugated synthetic peptide derived from Epstein-Barr virus Replication and transcription |  |
|   |  |  |
|   | activator  |  |
| Isotype:  |  |  |
| Isotype: Cross-Reactivity:                                    | activator  |  |
|   | activator IgG  |  |
| Cross-Reactivity:   | activator  IgG  Human, Virus   |  |
| Cross-Reactivity: Cross-Reactivity (Details):                 | activator  IgG  Human, Virus  HHV4tp2  |  |
| Cross-Reactivity:  Cross-Reactivity (Details):  Purification: | activator  IgG  Human, Virus  HHV4tp2  |  |

| Target Details      |  |  |
|---------------------|--|--|
| Target Type:        | Viral Protein  |  |
| Background:         | Synonyms: Replication and transcription activator, Rta, Immediate-early protein Rta, BRLF1 Background: Immediate-early transcription factor that controls the initiation of viral lytic gene expression and lytic reactivation from latency. Triggers lytic replication, and initiates a cellular senescence program in epithelial cells. Upregulates human DCR3/TNFRSF6B by directly binding to its receptor. |  |
| Gene ID:            | 3783727  |  |
| UniProt:            | P03209   |  |
| Application Details |  |  |
| Application Notes:  | WB 1:300-5000<br>IHC-P 1:200-400   |  |
| Restrictions:       | For Research Use only  |  |
| Handling            |  |  |
| Format:             | Liquid   |  |
| Concentration:      | 1 μg/μL  |  |
| Buffer:             | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.   |  |

This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

ProClin

-20 °C

12 months

handled by trained staff only.

Store at -20°C for 12 months.

Preservative:

Storage:

Expiry Date:

Precaution of Use:

Storage Comment: