antibodies .- online.com







anti-BTG3 antibody (AA 1-100) (HRP)



Overview

| Quantity: | 100 μL |
|----------------------|---|
| Target: | BTG3 |
| Binding Specificity: | AA 1-100 |
| Reactivity: | Human, Mouse, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This BTG3 antibody is conjugated to HRP |
| Application: | Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human BTG3 |
|-----------------------|--|
| Isotype: | IgG |
| Cross-Reactivity: | Human, Mouse, Rat |
| Predicted Reactivity: | Dog,Cow,Sheep,Pig,Horse |
| Purification: | Purified by Protein A. |

Target Details

| Target: | BTG3 |
|-------------------|----------------------|
| Alternative Name: | BTG3 (BTG3 Products) |

Target Details

| rarget betails | |
|---------------------|---|
| Background: | Synonyms: Abundant in neuroepithelium area protein, ANA, BTG family member 3, Btg3, BTG3_HUMAN, Protein BTG3, Protein Tob5, Protein BTG3. Background: The protein encoded by this gene is a member of the BTG/Tob family. This family has structurally related proteins that appear to have antiproliferative properties. This encoded protein might play a role in neurogenesis in the central nervous system. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Aug 2008]. |
| Gene ID: | 10950 |
| UniProt: | Q14201 |
| Application Details | |
| Application Notes: | WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500 |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Handling Advice: | Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date: | 12 months |