# antibodies -online.com





## anti-CACNA1A antibody (AA 1401-1500) (AbBy Fluor® 350)



Go to Product page

$\sim$					
	1/6	⊃r	\/I	$\triangle$	٨/

Quantity:	100 μL	
Target:	CACNA1A	
Binding Specificity:	AA 1401-1500	
Reactivity:	Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CACNA1A antibody is conjugated to AbBy Fluor® 350	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human CACNA1A
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human,Mouse,Cow,Pig,Rabbit
Purification:	Purified by Protein A.

### Target Details

Target:	CACNA1A
Alternative Name:	CACNA1A/Cav2.1 (CACNA1A Products)

Precaution of Use:

Target Details		
Background:	Synonyms: APCA, BI, Brain calcium channel 1, Brain calcium channel I, Cach4, Cacn3, Cacna1a,	
	Cacnl1a4, Calcium channel alpha 1A subunit, Calcium channel L type alpha 1 polypeptide,	
	Calcium channel L type alpha-1 polypeptide isoform 4, Calcium channel voltage dependent, P/Q	
	type alpha 1A subunit, CAV2.1, EA2, FHM, HPCA, MHP, MHP1, RAT brain class A, RBA-I, SCA6,	
	Voltage-dependent P/Q-type calcium channel alpha-1A subunit, Voltage-gated calcium channel	
	alpha subunit Cav2.1, CAC1A_HUMAN.	
	Background: Cav2.1 is a voltage-sensitive calcium channels (VSCC) which belongs to the	
	calcium channel alpha-1 subunit family. Cav2.1 mediates the entry of calcium ions into	
	excitable cells and is also involved in a variety of calcium-dependent processes, including	
	muscle contraction, hormone or neurotransmitter release, gene expression, cell motility, cell	
	division and cell death. Cav2.1 (isoform alpha-1A) gives rise to P and/or Q-type calcium	
	currents. Voltage-dependent calcium channels are multisubunit complexes, consisting of alpha-	
	1, alpha-2, beta and delta subunits in a 1:1:1:1 ratio. The channel activity is directed by the pore-	
	forming and voltage-sensitive alpha-1 subunit. In many cases, this subunit is sufficient to	
	generate voltage-sensitive calcium channel activity. The auxiliary subunits beta and alpha-	
	2/delta linked by a disulfide bridge regulate the channel activity.	
Gene ID:	773	
Pathways:	Regulation of Hormone Metabolic Process, Regulation of Hormone Biosynthetic Process	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	

This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

### Handling

	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months