antibodies.com

## Datasheet for ABIN886090 anti-CACNA1A antibody (AA 1401-1500) (Alexa Fluor 647)



## Overview

Quantity:	100 µL	
Target:	CACNA1A	
Binding Specificity:	AA 1401-1500	
Reactivity:	Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CACNA1A antibody is conjugated to Alexa Fluor 647	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CACNA1A	
lsotype:	lgG	
Cross-Reactivity:	Rat	
Predicted Reactivity:	Human,Mouse,Cow,Pig,Rabbit	
Purification:	Purified by Protein A.	
Target Details		
Taurati	04001414	

Target:	CACNA1A
Alternative Name:	CACNA1A/Cav2.1 (CACNA1A Products)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN886090 | 03/08/2024 | Copyright antibodies-online. All rights reserved.

Target Details	
Background:	Synonyms: APCA, BI, Brain calcium channel 1, Brain calcium channel I, Cach4, Cacn3, Cacna1a,
	Cacnl1a4, Calcium channel alpha 1A subunit, Calcium channel L type alpha 1 polypeptide,
	Calcium channel L type alpha-1 polypeptide isoform 4, Calcium channel voltage dependent, P/Q
	type alpha 1A subunit, CAV2.1, EA2, FHM, HPCA, MHP, MHP1, RAT brain class A, RBA-I, SCA6,
	Voltage-dependent P/Q-type calcium channel alpha-1A subunit, Voltage-gated calcium channel
	alpha subunit Cav2.1, CAC1A_HUMAN.
	Background: Cav2.1 is a voltage-sensitive calcium channels (VSCC) which belongs to the
	calcium channel alpha-1 subunit family. Cav2.1 mediates the entry of calcium ions into
	excitable cells and is also involved in a variety of calcium-dependent processes, including
	muscle contraction, hormone or neurotransmitter release, gene expression, cell motility, cell
	division and cell death. Cav2.1 (isoform alpha-1A) gives rise to P and/or Q-type calcium
	currents. Voltage-dependent calcium channels are multisubunit complexes, consisting of alpha-
	1, alpha-2, beta and delta subunits in a 1:1:1:1 ratio. The channel activity is directed by the pore-
	forming and voltage-sensitive alpha-1 subunit. In many cases, this subunit is sufficient to
	generate voltage-sensitive calcium channel activity. The auxiliary subunits beta and alpha-
	2/delta linked by a disulfide bridge regulate the channel activity.
Gene ID:	773
Pathways:	Regulation of Hormone Metabolic Process, Regulation of Hormone Biosynthetic Process
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN886090 | 03/08/2024 | Copyright antibodies-online. All rights reserved.

lond	lina
land	

	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months