antibodies -online.com





anti-NOD2 antibody (AA 251-350) (AbBy Fluor® 488)





Go to Product page

Overview

Quantity:	100 μL
Target:	NOD2
Binding Specificity:	AA 251-350
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NOD2 antibody is conjugated to AbBy Fluor® 488
Application:	Western Blotting (WB), Flow Cytometry (FACS)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CARD15
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	NOD2
Alternative Name:	NOD2 (NOD2 Products)
Background:	Synonyms: ACUG, Arthrocutaneouveal granulomatosis, BLAU, CARD15, Caspase recruitment

domain protein 15, Caspase recruitment domain-containing protein 15, IBD1, Inflammatory bowel disease protein 1, LRR containing protein, LRR-containing protein, Nod2, NoD2 protein, NoD2_HUMAN, NoD2B, nucleotide binding oligomerization domain 2, nucleotide-binding oligomerization domain 2, Nucleotide-binding oligomerization domain-containing protein 2, PSORAS1.

Background: Defects in NOD2 are the cause of sarcoidosis early-onset (EOS). EOS is a form of sarcoidosis manifesting in children younger than 4 years of age. Sarcoidosis is an idiopathic, systemic, inflammatory disease characterized by the formation of immune granulomas in involved organs. Granulomas predominantly invade the lungs and the lymphatic system, but also skin, liver, spleen, eyes and other organs may be involved. Early-onset sarcoidosis is quite rare and has a distinct triad of skin, joint and eye disorders, without apparent pulmonary involvement. Compared with an asymptomatic and sometimes naturally disappearing course of the disease in older children, early-onset sarcoidosis is progressive and in many cases causes severe complications, such as blindness, joint destruction and visceral involvement.

Gene ID:

UniProt: Q9HC29

64127

Pathways:

Activation of Innate immune Response, Cellular Response to Molecule of Bacterial Origin,
Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process,
Production of Molecular Mediator of Immune Response, Toll-Like Receptors Cascades,
Inflammasome

Application Details

Application Notes: FCM 1:20-100

Restrictions: For Research Use only


Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

Handling

	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months

Images



Flow Cytometry

Image 1. Raji probed with CARD15 Polyclonal Antibody, ALEXA FLUOR® 488 Conjugated (bs-7084R-A488) at 1:100 for 30 minutes compared to control cells (blue)and isotype control (orange).