antibodies -online.com





anti-CCDC138 antibody (AA 551-665) (Alexa Fluor 555)



Go to Product page

\sim					
()	VE	۲۱	/1	\triangle	Λ

Quantity:	100 μL	
Target:	CCDC138	
Binding Specificity:	AA 551-665	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CCDC138 antibody is conjugated to Alexa Fluor 555	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CCDC138	
Isotype:	IgG	
Cross-Reactivity:	Human	
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Pig,Horse,Rabbit	
Purification:	Purified by Protein A.	

Target Details

Target:	CCDC138	
Alternative Name:	CCDC138 (CCDC138 Products)	

Target Details

Background:	Synonyms: CCDC 138, CCDC138, Coiled coil domain containing 138, Coiled coil domain
	containing protein 138, coiled-coil domain containing 138, FLJ 32745, FLJ32745,
	CC138_HUMAN.
	Background: CCDC138, also known as FLJ32745, is a 685 amino acid protein expressed as two
	isoforms produced by alternative splicing. The gene that encodes CCDC138 maps to human
	chromosome 2q12.3. The second largest human chromosome, chromosome 2 consists of 237
	million bases, encodes over 1,400 genes and makes up approximately 8 % of the human
	genome. A number of genetic diseases are linked to genes on chromosome 2. Harlequin
	icthyosis, a rare and morbid skin deformity, is associated with mutations in the ABCA12 gene.
	The lipid metabolic disorder sitosterolemia is associated with ABCG5 and ABCG8. An extremely
	rare recessive genetic disorder, Alstr syndrome is due to mutations in the ALMS1 gene.
	Interestingly, chromosome 2 contains what appears to be a vestigial second centromere and
	vestigial telomeres which gives credence to the hypothesis that human chromosome 2 is the
	result of an ancient fusion of two ancestral chromosomes seen in modern form today in apes.
Gene ID:	165055
Pathways:	BCR Signaling
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C

Handling

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months