



[Go to Product page](#)

Datasheet for ABIN887485

anti-CCDC18 antibody (AA 55-160) (Alexa Fluor 488)

Overview

Quantity:	100 µL
Target:	CCDC18
Binding Specificity:	AA 55-160
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CCDC18 antibody is conjugated to Alexa Fluor 488
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CCDC18
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	CCDC18
Alternative Name:	CCDC18 (CCDC18 Products)
Background:	Synonyms: CCD18_HUMAN, Ccdc18, Coiled-coil domain-containing protein 18, Sarcoma

Target Details

antigen NY-SAR-24.

Background: CCDC18, also known as NY-SAR-41 or dJ717I23.1, is a 1,454 amino acid protein expressed as two isoforms and encoded by a gene mapping to human chromosome 1.

Chromosome 1 is the largest human chromosome spanning about 260 million base pairs and making up 8 % of the human genome. There are about 3,000 genes on chromosome 1, and considering the great number of genes there are also a large number of diseases associated with chromosome 1. Notably, the rare aging disease Hutchinson-Gilford progeria is associated with the LMNA gene which encodes lamin A. When defective, the LMNA gene product can build up in the nucleus and cause characteristic nuclear blebs. The mechanism of rapidly enhanced aging is unclear and is a topic of continuing exploration. The MUTYH gene is located on chromosome 1 and is partially responsible for familial adenomatous polyposis. Stickler syndrome, Parkinsons, Gaucher disease and Usher syndrome are also associated with chromosome 1. A breakpoint has been identified in 1q which disrupts the DISC1 gene and is linked to schizophrenia. Aberrations in chromosome 1 are found in a variety of cancers including head and neck cancer, malignant melanoma and multiple myeloma.

Gene ID: 343099

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Handling

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months