Datasheet for ABIN887842 anti-CCDC50 antibody (AA 251-306) (Biotin)

antibodies.com



Overview

Quantity:	100 µL
Target:	CCDC50
Binding Specificity:	AA 251-306
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CCDC50 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CCDC50
Isotype:	lgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat
Purification:	Purified by Protein A.
Target Details	
Target:	CCDC50
Alternative Name:	CCDC50 (CCDC50 Products)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN887842 | 03/06/2024 | Copyright antibodies-online. All rights reserved.

Target Details

Synonyms: C3orf6, C3orf6 YMER, CCD50_HUMAN, CCDC 50, Ccdc50, Chromosome 3 open
reading frame 6, Coiled coil domain containing 50, Coiled coil domain containing protein 50,
Coiled-coil domain-containing protein 50, Protein Ymer, Ymer, Ymer protein.
Background: This gene encodes a soluble, cytoplasmic, tyrosine-phosphorylated protein with
multiple ubiquitin-interacting domains. Mutations in this gene cause nonsyndromic, postlingual,
progressive sensorineural DFNA44 hearing loss. In mouse, the protein is expressed in the inner
ear during development and postnatal maturation and associates with microtubule-based
structures. This protein may also function as a negative regulator of NF-kB signaling and as an
effector of epidermal growth factor (EGF)-mediated cell signaling. Alternative splicing results in
multiple transcript variants encoding distinct isoforms. [provided by RefSeq, Oct 2008].

Gene ID:

152137

Application Details

Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months