antibodies -online.com





Datasheet for ABIN888066

anti-CCDC69 antibody (AA 41-140) (Alexa Fluor 488)

Go to Product page

()	1 /	0	rv	/ 1 /	71	Α.
	1//	-	1 \/	16		1/1/
\sim	v	\sim	1 V	١,	_	v v

Quantity:	100 μL	
Target:	CCDC69	
Binding Specificity:	AA 41-140	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CCDC69 antibody is conjugated to Alexa Fluor 488	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CCDC69
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Dog,Cow,Sheep,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	CCDC69
Alternative Name:	CCDC69 (CCDC69 Products)

Target Details

Expiry Date:

Background:	Synonyms: CCD69_HUMAN, ccdc69, Coiled coil domain containing 69, Coiled-coil domain-
	containing protein 69.
	Background: The coiled-coil domain is a structural motif found in proteins that are involved in a
	diverse array of biological functions such as the regulation of gene expression, cell division,
	membrane fusion and drug extrusion and delivery. CCDC69 (Coiled-coil domain-containing
	protein 69) is a 296 amino acid protein that is encoded by a gene which maps to human
	chromosome 5, which contains 181 million base pairs and comprises nearly 6 % of the human
	genome. Chromosome 5 is associated with Cockayne syndrome through the ERCC8 gene and
	familial adenomatous polyposis through the adenomatous polyposis coli (APC) tumor
	suppressor gene. Treacher Collins syndrome is also chromosome 5-associated and is caused
	by insertions or deletions within the TCOF1 gene. Deletion of the p arm of chromosome 5 leads
	to Cri du chat syndrome, while deletion of the q arm or of chromosome 5 altogether is common
	in therapy-related acute myelogenous leukemias and myelodysplastic syndrome.
Gene ID:	26112
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

12 months