antibodies -online.com





anti-CCDC98 antibody (AA 51-150) (Alexa Fluor 555)



Go to Product page

()	11/0	K\ /	iew	1
	\cup	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL	
Target:	CCDC98 (FAM175A)	
Binding Specificity:	AA 51-150	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CCDC98 antibody is conjugated to Alexa Fluor 555	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CCDC98	
Isotype:	IgG	
Cross-Reactivity:	Mouse	
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Pig,Horse,Chicken	
Purification:	Purified by Protein A.	

Target Details

Target:	CCDC98 (FAM175A)
Alternative Name:	CCDC98 (FAM175A Products)

Target Details

Background:	Synonyms: ABRA1, CCDC98, BRCA1-A complex subunit Abraxas, Coiled-coil domain-containing	
	protein 98, Protein FAM175A, FAM175A, UNQ496/PRO1013	
	Background: Involved in DNA damage response and double-strand break (DSB) repair.	
	Component of the BRCA1-A complex, acting as a central scaffold protein that assembles the	
	various components of the complex and mediates the recruitment of BRCA1. The BRCA1-A	
	complex specifically recognizes 'Lys-63'-linked ubiquitinated histones H2A and H2AX at DNA	
	lesion sites, leading to target the BRCA1-BARD1 heterodimer to sites of DNA damage at DSBs.	
	This complex also possesses deubiquitinase activity that specifically removes 'Lys-63'-linked	
	ubiquitin on histones H2A and H2AX.	
Gene ID:	84142	
UniProt:	Q6UWZ7	
Pathways:	DNA Damage Repair, Positive Regulation of Response to DNA Damage Stimulus	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	