antibodies -online.com





anti-CCL19 antibody (AA 26-108) (AbBy Fluor® 488)



Go to Product page

\sim					
()	VE	۲۱	/1	\triangle	Λ

Quantity:	100 μL	
Target:	CCL19	
Binding Specificity:	AA 26-108	
Reactivity:	Human, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CCL19 antibody is conjugated to AbBy Fluor® 488	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from mouse CCL19
Isotype:	IgG
Cross-Reactivity:	Human, Rat
Predicted Reactivity:	Mouse,Pig
Purification:	Purified by Protein A.

Target Details

Target:	CCL19	
Alternative Name:	Ccl19/Mip-3 beta (CCL19 Products)	

Target Details

Synonyms: ELC, CKb11, MIP3B, Scya19, exodus-3, C-C motif chemokine 19, Epstein-Barr virus-induced molecule 1 ligand chemokine, EBI1 ligand chemokine, Small-inducible cytokine A19,		
Ccl19		
Background: Strongly chemotactic for naive (L-selectinhi) CD4 T-cells and for CD8 T-cells and		
weakly attractive for resting B-cells and memory (L-selectinlo) CD4 T-cells. May play a role in		
promoting encounters between recirculating T-cells and dendritic cells and in the migration of		
activated B-cells into the T-zone of secondary lymphoid tissues. Binds to chemokine receptor		
CCR7. Binds to atypical chemokine receptor ACKR4 and mediates the recruitment of beta-		
arrestin (ARRB1/2) to ACKR4.		
24047		
070460		
Positive Regulation of Immune Effector Process		
IF(IHC-P) 1:50-200		
IF(IHC-F) 1:50-200		
IF(ICC) 1:50-200		
For Research Use only		
Liquid		
1 μg/μL		
Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and		
50 % Glycerol.		
ProClin		
This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be		
handled by trained staff only.		
-20 °C		
Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		
12 months		