# antibodies .- online.com







## anti-Septin 5 antibody (AA 21-120) (Cy3)



( )	1/0	r\ /1	014	
( )	ve	I V I	-v	V

Quantity:	100 μL	
Target:	Septin 5 (SEPT5)	
Binding Specificity:	AA 21-120	
Reactivity:	Human, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Septin 5 antibody is conjugated to Cy3	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human CDCREL/SEPT5	
Isotype:	IgG	
Cross-Reactivity:	Human, Rat	
Predicted Reactivity:	Mouse,Dog,Cow,Pig,Horse	
Purification:	Purified by Protein A.	

### Target Details

Target:	Septin 5 (SEPT5)
Alternative Name: Septin5 (SEPT5 Products)	

### Target Details

Background:	Synonyms: 5 Sep, CDCrel 1, CDCREL, CDCrel-1, CDCREL1, Cell division control related protein 1,	
	Cell division control-related protein 1, H5 antibody HCDCREL 1, Peanut like 1, Peanut like 1	
	homolog, Peanut like protein 1, Peanut-like protein 1, Platelet glycoprotein lb beta chain,	
	PNUTL1, SEPT5, SEPT-5, SEPT 5, SEPT5_HUMAN, Septin 5, Septin5, Septin-5.	
	Background: This gene is a member of the septin gene family of nucleotide binding proteins,	
	originally described in yeast as cell division cycle regulatory proteins. Septins are highly	
	conserved in yeast, Drosophila, and mouse and appear to regulate cytoskeletal organization.	
	Disruption of septin function disturbs cytokinesis and results in large multinucleate or polyploid	
	cells. This gene is mapped to 22q11, the region frequently deleted in DiGeorge and	
	velocardiofacial syndromes. A translocation involving the MLL gene and this gene has also	
	been reported in patients with acute myeloid leukemia. Alternative splicing results in multiple	
	transcript variants. The presence of a non-consensus polyA signal (AACAAT) in this gene also	
	results in read-through transcription into the downstream neighboring gene (GP1BB, platelet	
	glycoprotein lb), whereby larger, non-coding transcripts are produced. [provided by RefSeq, Dec	
	2010].	
Gene ID:	5413	
UniProt:	Q99719	
Pathways:	Synaptic Vesicle Exocytosis	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	

### Handling

	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months