

Datasheet for ABIN889717

anti-Septin 5 antibody (AA 21-120) (Cy5)



()	11 /	er	~\ /	10	1 A /
	1 / /	-1	١/	-	1///
\sim	٧.	\sim 1	v	\sim	· v v

Overview	
Quantity:	100 μL
Target:	Septin 5 (SEPT5)
Binding Specificity:	AA 21-120
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Septin 5 antibody is conjugated to Cy5
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human CDCREL/SEPT5
Isotype:	IgG
Cross-Reactivity:	Human, Rat
Predicted Reactivity:	Mouse,Dog,Cow,Pig,Horse
Purification:	Purified by Protein A.
Target Details	
Target:	Septin 5 (SEPT5)
Alternative Name:	Septin5 (SEPT5 Products)

Target Details

Precaution of Use:

ranger betane			
Background:	Synonyms: 5 Sep, CDCrel 1, CDCREL, CDCrel-1, CDCREL1, Cell division control related protein 1,		
	Cell division control-related protein 1, H5 antibody HCDCREL 1, Peanut like 1, Peanut like 1		
	homolog, Peanut like protein 1, Peanut-like protein 1, Platelet glycoprotein lb beta chain,		
	PNUTL1, SEPT5, SEPT-5, SEPT 5, SEPT5_HUMAN, Septin 5, Septin5, Septin-5.		
	Background: This gene is a member of the septin gene family of nucleotide binding proteins,		
	originally described in yeast as cell division cycle regulatory proteins. Septins are highly		
	conserved in yeast, Drosophila, and mouse and appear to regulate cytoskeletal organization.		
	Disruption of septin function disturbs cytokinesis and results in large multinucleate or polyploid		
	cells. This gene is mapped to 22q11, the region frequently deleted in DiGeorge and		
	velocardiofacial syndromes. A translocation involving the MLL gene and this gene has also		
	been reported in patients with acute myeloid leukemia. Alternative splicing results in multiple		
	transcript variants. The presence of a non-consensus polyA signal (AACAAT) in this gene also		
	results in read-through transcription into the downstream neighboring gene (GP1BB, platelet		
	glycoprotein lb), whereby larger, non-coding transcripts are produced. [provided by RefSeq, Dec		
	2010].		
Gene ID:	5413		
UniProt:	Q99719		
Pathways:	Synaptic Vesicle Exocytosis		
Application Details			
Application Notes:	IF(IHC-P) 1:50-200		
	IF(IHC-F) 1:50-200		
	IF(ICC) 1:50-200		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and		
	50 % Glycerol.		
Preservative:	ProClin		

This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

Handling

	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months