

Datasheet for ABIN889724

anti-Septin 5 antibody (AA 21-120) (Cy5.5)



0				

Quantity:	100 μL
Target:	Septin 5 (SEPT5)
Binding Specificity:	AA 21-120
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Septin 5 antibody is conjugated to Cy5.5
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	

Immunogen:	KLH conjugated synthetic peptide derived from human CDCREL/SEPT5
Isotype:	IgG
Cross-Reactivity:	Human, Rat
Predicted Reactivity:	Mouse,Dog,Cow,Pig,Horse
Purification:	Purified by Protein A.

Target Details

Target:	Septin 5 (SEPT5)
Alternative Name:	Septin5 (SEPT5 Products)

Target Details

Precaution of Use:

Synonyms: 5 Sep, CDCrel 1, CDCREL, CDCrel-1, CDCREL1, Cell division control related protein 1,
Cell division control-related protein 1, H5 antibody HCDCREL 1, Peanut like 1, Peanut like 1
homolog, Peanut like protein 1, Peanut-like protein 1, Platelet glycoprotein lb beta chain,
PNUTL1, SEPT5, SEPT-5, SEPT 5, SEPT5_HUMAN, Septin 5, Septin5, Septin-5.
Background: This gene is a member of the septin gene family of nucleotide binding proteins,
originally described in yeast as cell division cycle regulatory proteins. Septins are highly
conserved in yeast, Drosophila, and mouse and appear to regulate cytoskeletal organization.
Disruption of septin function disturbs cytokinesis and results in large multinucleate or polyploid
cells. This gene is mapped to 22q11, the region frequently deleted in DiGeorge and
velocardiofacial syndromes. A translocation involving the MLL gene and this gene has also
been reported in patients with acute myeloid leukemia. Alternative splicing results in multiple
transcript variants. The presence of a non-consensus polyA signal (AACAAT) in this gene also
results in read-through transcription into the downstream neighboring gene (GP1BB, platelet
glycoprotein lb), whereby larger, non-coding transcripts are produced. [provided by RefSeq, Dec
2010].
5413
Q99719
Synaptic Vesicle Exocytosis
IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200
For Research Use only
Liquid
1 μg/μL
Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
ProClin

This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

Handling

	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months