# antibodies -online.com





## anti-CDK2AP2 antibody (AA 75-126) (HRP)



Go to Product page

$\sim$					
()	VE	۲۱	/1	$\triangle$	Λ

Quantity:	100 μL
Target:	CDK2AP2
Binding Specificity:	AA 75-126
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CDK2AP2 antibody is conjugated to HRP
Application:	ELISA, Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human CDK2AP2
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Chicken
Purification:	Purified by Protein A.

#### **Target Details**

Target:	CDK2AP2
Alternative Name:	CDK2AP2 (CDK2AP2 Products)
Background:	Synonyms: CDK2 associated protein 2, CDK2-associated protein 2, CDK2AP2, CDKA2_HUMAN,

2, DOC 1 related protein, DOC-1-related protein, DOC-1R, DOC1R, FLJ10636, p14, tumor suppressor deleted in oral cancer related 1.  Background: This gene encodes a protein that interacts with cyclin-dependent kinase 2 associated protein 1. Pseudogenes associated with this gene are located on chromosomes 7 and 9. Alternatively spliced transcript variants have been observed for this gene. [provided by RefSeq. Dec 2012].	Cyclin dependent kinase 2 associated protein 2, Cyclin-dependent kinase 2-associated protein
Background: This gene encodes a protein that interacts with cyclin-dependent kinase 2 associated protein 1. Pseudogenes associated with this gene are located on chromosomes 7 and 9. Alternatively spliced transcript variants have been observed for this gene. [provided by	2, DOC 1 related protein, DOC-1-related protein, DOC-1R, DOC1R, FLJ10636, p14, tumor
associated protein 1. Pseudogenes associated with this gene are located on chromosomes 7 and 9. Alternatively spliced transcript variants have been observed for this gene. [provided by	suppressor deleted in oral cancer related 1.
and 9. Alternatively spliced transcript variants have been observed for this gene. [provided by	Background: This gene encodes a protein that interacts with cyclin-dependent kinase 2
	associated protein 1. Pseudogenes associated with this gene are located on chromosomes 7
RefSea, Dec 2012].	and 9. Alternatively spliced transcript variants have been observed for this gene. [provided by
The state of the s	RefSeq, Dec 2012].

Gene ID: 10263

Pathways: PI3K-Akt Signaling, Cell Division Cycle, Mitotic G1-G1/S Phases, DNA Replication, M Phase,

Synthesis of DNA

#### **Application Details**

Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only

### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months