



[Go to Product page](#)

Datasheet for ABIN890174
anti-Cep76 antibody (AA 451-550) (FITC)

Overview

| | |
|----------------------|---|
| Quantity: | 100 µL |
| Target: | Cep76 |
| Binding Specificity: | AA 451-550 |
| Reactivity: | Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This Cep76 antibody is conjugated to FITC |
| Application: | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| | |
|-----------------------|---|
| Immunogen: | KLH conjugated synthetic peptide derived from human CEP76 |
| Isotype: | IgG |
| Cross-Reactivity: | Rat |
| Predicted Reactivity: | Human,Mouse,Pig,Horse,Rabbit |
| Purification: | Purified by Protein A. |

Target Details

| | |
|-------------------|--|
| Target: | Cep76 |
| Alternative Name: | CEP76 (Cep76 Products) |

Target Details

Background: Synonyms: C18orf9, Centrosomal protein 76 kDa, Centrosomal protein of 76 kDa, CEP 76, Cep76, CEP76_HUMAN, Chromosome 18 open reading frame 9, FLJ12542, HsT1705.
Background: Centrosomal protein involved in regulation of centriole duplication. Required to limit centriole duplication to once per cell cycle by preventing centriole reduplication.

Gene ID: 79959

Pathways: [M Phase](#)

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months
