# antibodies -online.com





## anti-RCBTB1 antibody (AA 481-531) (Alexa Fluor 647)



Go to Product page

| ( ) | 11/0   | K\ /   | iew                 | 1 |
|-----|--------|--------|---------------------|---|
|     | $\cup$ | 'I V/I | $\square \vee \vee$ | ı |
|     |        |        |                     |   |

| Quantity:            | 100 μL  |  |
|----------------------|---|--|
| Target:              | RCBTB1  |  |
| Binding Specificity: | AA 481-531  |  |
| Reactivity:          | Rat   |  |
| Host:                | Rabbit  |  |
| Clonality:           | Polyclonal  |  |
| Conjugate:           | This RCBTB1 antibody is conjugated to Alexa Fluor 647   |  |
| Application:         | Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc)) |  |

#### **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human CLLD7 |
|-----------------------|---|
| Isotype:              | IgG   |
| Cross-Reactivity:     | Rat   |
| Predicted Reactivity: | Human,Mouse,Dog,Cow,Sheep,Pig,Horse,Chicken               |
| Purification:         | Purified by Protein A.                                    |

#### **Target Details**

| Target:           | RCBTB1                  |
|-------------------|-------------------------|
| Alternative Name: | CLLD7 (RCBTB1 Products) |

### Target Details

| 9                   |  |  |
|---------------------|--|--|
| Background:         | Synonyms: RCC1 and BTB domain-containing protein 1, Chronic lymphocytic leukemia deletion                  |  |
|                     | region gene 7 protein, CLL deletion region gene 7 protein, Regulator of chromosome                         |  |
|                     | condensation and BTB domain-containing protein 1, RCBT1_HUMAN.   |  |
|                     | Background: This gene encodes a protein with an N-terminal RCC1 domain and a C-terminal                    |  |
|                     | BTB (broad complex, tramtrack and bric-a-brac) domain. In rat, over-expression of this gene in             |  |
|                     | vascular smooth muscle cells induced cellular hypertrophy. In rat, the C-terminus of RCBTB1                |  |
|                     | interacts with the angiotensin II receptor-1A. In humans, this gene maps to a region of                    |  |
|                     | chromosome 13q that is frequently deleted in B-cell chronic lymphocytic leukemia and other                 |  |
|                     | lymphoid malignancies. [provided by RefSeq, Jul 2008].   |  |
| Gene ID:            | 55213  |  |
| Application Details |  |  |
| Application Notes:  | IF(IHC-P) 1:50-200   |  |
|                     | IF(IHC-F) 1:50-200   |  |
|                     | IF(ICC) 1:50-200   |  |
| Restrictions:       | For Research Use only  |  |
| Handling            |  |  |
| Format:             | Liquid   |  |
| Concentration:      | 1 μg/μL  |  |
| Buffer:             | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |  |
| Preservative:       | ProClin  |  |
| Precaution of Use:  | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be                        |  |
|                     | handled by trained staff only.   |  |
| Storage:            | -20 °C   |  |
| Storage Comment:    | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                          |  |
| Expiry Date:        | 12 months  |  |
|                     |  |  |