antibodies .- online.com





anti-CNGA2 antibody (AA 155-250) (Alexa Fluor 647)



()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μL	
Target:	CNGA2	
Binding Specificity:	AA 155-250	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CNGA2 antibody is conjugated to Alexa Fluor 647	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CNGA2	
Isotype:	IgG	
Specificity:	This antibody may have a secondary reaction to Dapk1 due to a 61 % contiguous sequence in the immunogen region.	
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Horse, Chicken, Rabbit	
Purification:	Purified by Protein A.	

Target Details

Target:	CNGA2

Target Details

rarget betails		
Alternative Name:	Cnga2 (CNGA2 Products)	
Background:	Synonyms: CNCA, CNG2, CNCA1, OCNC1, OCNCa, OCNCALPHA, Cyclic nucleotide-gated	
	olfactory channel, Cyclic nucleotide-gated cation channel 2, Cyclic nucleotide-gated channel	
	alpha-2, CNG channel alpha-2, CNG-2, CNGA2, CNCG2	
	Background: Odorant signal transduction is probably mediated by a G-protein coupled cascade	
	using cAMP as second messenger. The olfactory channel can be shown to be activated by	
	cyclic nucleotides which leads to a depolarization of olfactory sensory neurons.	
Gene ID:	1260	
UniProt:	Q16280	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	