

[Go to Product page](#)

Datasheet for ABIN892066

anti-COX7A2 antibody (AA 21-83) (Alexa Fluor 488)

Overview

Quantity:	100 µL
Target:	COX7A2
Binding Specificity:	AA 21-83
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This COX7A2 antibody is conjugated to Alexa Fluor 488
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human COX7A2
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	COX7A2
Alternative Name:	COX7A2 (COX7A2 Products)
Background:	Synonyms: Mitochondrion. Cytochrome c oxidase subunit 7A2, mitochondrial, COX7a related

Target Details

protein, Cytochrome c oxidase subunit VIIa-L, Cytochrome c oxidase subunit VIIa-liver/heart, COX7a-related protein, COX7AR, COX7RP, Cytochrome c oxidase subunit VIIa related protein, mitochondrial, Cytochrome c oxidase subunit VIIaL, Cytochrome c oxidase subunit VIIa-L, Cytochrome c oxidase subunit VIIa-related protein, mitochondrial [Precursor], EB1, Estrogen receptor binding CpG island, SIG81, CX7A2_HUMAN.

Background: Cytochrome c oxidase, the terminal component of the mitochondrial respiratory chain, catalyzes the electron transfer from reduced cytochrome c to oxygen. This component is a heteromeric complex consisting of three catalytic subunits encoded by mitochondrial genes, and multiple structural subunits encoded by nuclear genes. The mitochondrially-encoded subunits function in electron transfer, while the nuclear-encoded subunits may function in the regulation and assembly of the complex. This nuclear gene encodes polypeptide 2 (liver isoform) of subunit VIIa, with this polypeptide being present in both muscle and non-muscle tissues. In addition to polypeptide 2, subunit VIIa includes polypeptide 1 (muscle isoform), which is present only in muscle tissues, and a related protein, which is present in all tissues. Alternative splicing results in multiple transcript variants. Related pseudogenes have been identified on chromosomes 4 and 14. [provided by RefSeq, Oct 2009]

Gene ID: 1347

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling

Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months