antibodies

## Datasheet for ABIN892324 anti-CRLF2 antibody (AA 81-180) (AbBy Fluor® 350)



Overview

Quantity:	100 µL	
Target:	CRLF2	
Binding Specificity:	AA 81-180	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CRLF2 antibody is conjugated to AbBy Fluor® 350	
Application:	Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))	

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CRLF2	
Isotype:	IgG	
Predicted Reactivity:	Human	
Purification:	Purified by Protein A.	

## Target Details

Target:	CRLF2
Alternative Name:	Crlf2 (CRLF2 Products)
Background:	Synonyms: TSLPR, CRLF2, CRL 2, CRL2, CRLF 2, CRLF2Y, Cytokine receptor CRL2 precusor,

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN892324 | 06/07/2024 | Copyright antibodies-online. All rights reserved.

	Cytokine receptor like 2, Cytokine receptor like factor 2, Cytokine receptor like factor 2
	precursor, IL XR, ILXR, Thymic stromal derived lymphopoietin receptor, Thymic stromal
	lymphopoietin protein receptor, Thymic stromal lymphopoietin receptor, TSLP receptor, TSLPR.
	Background: TSLPR is a type I membrane receptor that forms a functional heterodimeric
	complex with IL7R to bind TSLP. The TSLP R contains a WSXWS motif required for proper
	protein folding and a box1 motif important for association with the JAKs. TSLPR has a
	predicted molecular weight approximately 41 kD, and two further isoforms have been reported
	that are produced by alternative splicing. The TSLPR is expressed preferentially in myeloid cells
	including dendritic cells and activated monocytes, and is weakly expressed in T cells.
	Expression has also been reported in heart, skeletal muscle, and kidney tissues. TSLP binding
	to the heterodimeric functional receptor (TSLPR and IL7R) activates JAK2, STAT3 and STAT5
	to stimulate cell proliferation. Ligand receptor interactions haves been implicated in the
	development of the hematopoietic system, dendritic cell maturation, and the maintenance and
	polarization of human Th2 memory T cells in allergic diseases.
Gene ID:	64109

## Application Details

Format:	Liquid
Handling	
Restrictions:	For Research Use only
	IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Application Notes:	IF(IHC-P) 1:50-200

Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN892324 | 06/07/2024 | Copyright antibodies-online. All rights reserved.

1.1	(	1:
Н	land	ling
		3

Expiry Date:

12 months

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN892324 | 06/07/2024 | Copyright antibodies-online. All rights reserved.