antibodies -online.com





anti-CUBN antibody (AA 51-150) (Alexa Fluor 488)



Go to Product page

()	11/	IN	/ie	A .
	/ // 	۱ ات	/ (−	' \/\/

Quantity:	100 μL	
Target:	CUBN	
Binding Specificity:	AA 51-150	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CUBN antibody is conjugated to Alexa Fluor 488	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Cubilin	
Isotype:	IgG	
Cross-Reactivity:	Human	
Predicted Reactivity:	Mouse,Rat	
Purification:	Purified by Protein A.	

Target Details

Target:	CUBN
Alternative Name:	Cubilin (CUBN Products)

Target Details

9			
Background:	Synonyms: IFCR, MGA1, gp280, Cubilin, 460 kDa receptor, Intestinal intrinsic factor receptor,		
	Intrinsic factor-cobalamin receptor, Intrinsic factor-vitamin B12 receptor, CUBN		
	Background: Cotransporter which plays a role in lipoprotein, vitamin and iron metabolism, by		
	facilitating their uptake. Binds to ALB, MB, Kappa and lambda-light chains, TF, hemoglobin, GC,		
	SCGB1A1, APOA1, high density lipoprotein, and the GIF-cobalamin complex. The binding of all		
	ligands requires calcium. Serves as important transporter in several absorptive epithelia,		
	including intestine, renal proximal tubules and embryonic yolk sac. Interaction with LRP2		
	mediates its trafficking throughout vesicles and facilitates the uptake of specific ligands like		
	GC, hemoglobin, ALB, TF and SCGB1A1. Interaction with AMN controls its trafficking to the		
	plasma membrane and facilitates endocytosis of ligands. May play an important role in the		
	development of the peri-implantation embryo through internalization of APOA1 and cholesterol.		
	Binds to LGALS3 at the maternal-fetal interface.		
Gene ID:	8029		
UniProt:	060494		
Pathways:	Metabolism of Steroid Hormones and Vitamin D, Lipid Metabolism		
Application Details			
Application Notes:	IF(IHC-P) 1:50-200		
	IF(IHC-F) 1:50-200		
	IF(ICC) 1:50-200		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and		
	50 % Glycerol.		
Preservative:	ProClin		
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be		
	handled by trained staff only.		
Storage:	-20 °C		

Handling

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months