

[Go to Product page](#)

Datasheet for ABIN893046

anti-Cyclin T2 antibody (AA 185-270) (Cy5)

Overview

| | |
|----------------------|---|
| Quantity: | 100 µL |
| Target: | Cyclin T2 (CCNT2) |
| Binding Specificity: | AA 185-270 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This Cyclin T2 antibody is conjugated to Cy5 |
| Application: | Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc)) |

Product Details

| | |
|-----------------------|---|
| Immunogen: | KLH conjugated synthetic peptide derived from human Cyclin T2 |
| Isotype: | IgG |
| Predicted Reactivity: | Human, Mouse, Rat, Cow, Pig, Horse, Chicken, Rabbit |
| Purification: | Purified by Protein A. |

Target Details

| | |
|-------------------|--|
| Target: | Cyclin T2 (CCNT2) |
| Alternative Name: | Cyclin T2 (CCNT2 Products) |
| Background: | Synonyms: CCNT2, CCNT2_HUMAN, Cyclin-T2, CycT2, FLJ90560, MGC134840. |

Target Details

Background: Regulatory subunit of the cyclin-dependent kinase pair (CDK9/cyclin T) complex, also called positive transcription elongation factor B (P-TEFB), which is proposed to facilitate the transition from abortive to production elongation by phosphorylating the CTD (carboxy-terminal domain) of the large subunit of RNA polymerase II (RNAP II).

Gene ID: 905

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months