antibodies -online.com





anti-Cytochrome P450, Family 2, Subfamily J, Polypeptide 3 (CYP2J3) (AA 401-502) antibody (FITC)



Go to Product page

()	۱ ۱	\cap	r	/1	\cap	۱ ۸	1
0	'V	ㄷ	I١	νı	\Box	V	۷

Quantity:	100 μL
Target:	Cytochrome P450, Family 2, Subfamily J, Polypeptide 3 (CYP2J3)
Binding Specificity:	AA 401-502
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	FITC
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Cyp2-j3	
Isotype:	IgG	
Cross-Reactivity:	Mouse, Rat	
Purification:	Purified by Protein A.	

Target Details

Target:	Cytochrome P450, Family 2, Subfamily J, Polypeptide 3 (CYP2J3)	
Alternative Name:	Cyp2-j3 (CYP2J3 Products)	
Background:	Synonyms: CPJ2, Cytochrome P450 2J2, Arachidonic acid epoxygenase, CYPIIJ2, CYP2J2	

Target Details

	Background: This enzyme metabolizes arachidonic acid predominantly via a NADPH-dependent	
	olefin epoxidation to all four regioisomeric cis-epoxyeicosatrienoic acids. One of the	
	predominant enzymes responsible for the epoxidation of endogenous cardiac arachidonic acid	
	pools.	
Molecular Weight:	55kDa	
Gene ID:	1573	
UniProt:	P51589	

Application Details

Restrictions:	For Research Use only
	IF(ICC) 1:50-200
	IF(IHC-F) 1:50-200
Application Notes:	IF(IHC-P) 1:50-200

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months