antibodies.com

Datasheet for ABIN893541 anti-DBF4 antibody (AA 21-110) (Cy5)



0		
$() \times ic$	>r\/1	O111
Ove	-1 V I	
\sim		~

Quantity:	100 μL
Target:	DBF4
Binding Specificity:	AA 21-110
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DBF4 antibody is conjugated to Cy5
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human DBF4A
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Cow,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	DBF4
Alternative Name:	DBF4A (DBF4 Products)
Background:	Synonyms: Activator of S phase kinase, ASK, Chfon homolog A, DBF4, DBF4 type zinc finger

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN893541 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

Target Details

	containing protein 1, DBF4-type zinc finger-containing protein 1, DBF4A, DBF4A_HUMAN, Protein DBF4 homolog A, ZDBF1.
	Background: Regulatory subunit for CDC7 which activates its kinase activity thereby playing a central role in DNA replication and cell proliferation. Required for progression of S phase. The complex CDC7-DBF4A selectively phosphorylates MCM2 subunit at 'Ser-40' and 'Ser-53' and then is involved in regulating the initiation of DNA replication during cell cycle. Tissue specificity: Highly expressed in testis and thymus. Expressed also in most cancer cells lines.
Gene ID:	10926
Pathways:	Mitotic G1-G1/S Phases, DNA Replication
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months