# antibodies -online.com



Datasheet for ABIN893800

# anti-C-Type Lectin Domain Family 6, Member A (CLEC6A) (AA 110-209) antibody (AbBy Fluor® 350)



( )	11/	IN	/ie	A .
	/ // <del> </del>	۱ ات	/   (−	' \/\/

Quantity:	100 μL	
Target:	C-Type Lectin Domain Family 6, Member A (CLEC6A)	
Binding Specificity:	AA 110-209	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	AbBy Fluor® 350	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human Dectin-2
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Cow,Sheep,Horse
Purification:	Purified by Protein A.

# **Target Details**

Target:	C-Type Lectin Domain Family 6, Member A (CLEC6A)	
Alternative Name:	DECTIN 2 (CLEC6A Products)	
Background: Synonyms: C type lectin domain family 6 member A, C type lectin superfamily member 10, 0		

type lectin domain family 6 member A, C-type lectin superfamily member 10, CLC6A\_HUMAN, CLEC4N, Clec6a, CLECSF10, DC associated C type lectin 2, DC-associated C-type lectin 2, Dectin-2, Dectin-2, Dectin-2, Dendritic cell associated lectin 2, Dendritic cell-associated C-type lectin 2. Background: Dectin 2 is a type II transmembrane protein that is a member of the C type lectin superfamily. In mouse, dectin 2 is predominantly expressed on tissue macrophages and some dendritic cells. Significant expression of Dectin 2 has been reported on macrophages in the red pulp and marginal zones of the spleen, kupffer cells in the liver and alveolar macrophages in the lung. Peripheral blood monocytes express low levels of dectin 2 but transient up regulation of expression has been demonstrated on monocytes at sites of inflammation. The function of dectin 2 has not been fully determined but studies suggest a possible role in immune surveillance.

Gene ID:

93978

### **Application Details**

Application Notes: IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

## Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months