antibodies .- online.com





anti-Deltex Homolog 1 antibody (AA 351-450) (AbBy Fluor® 555)



Go to Product page

()	۱ ۱	\cap	r	/1	\cap	۱ ۸	1
0	'V	ㄷ	ı١	νı	\Box	V	۷

Quantity:	100 μL
Target:	Deltex Homolog 1 (DTX1)
Binding Specificity:	AA 351-450
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Deltex Homolog 1 antibody is conjugated to AbBy Fluor® 555
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Deltex-1
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Cow
Purification:	Purified by Protein A.

Target Details

Target:	Deltex Homolog 1 (DTX1)
Alternative Name:	Deltex 1 (DTX1 Products)

Target Details

Background:	Synonyms: hDx-1, E3 ubiquitin-protein ligase DTX1, Protein deltex-1, Deltex1, hDTX1, DTX1	
	Background: Functions as a ubiquitin ligase protein in vivo, mediating ubiquitination and	
	promoting degradation of MEKK1, suggesting that it may regulate the Notch pathway via some	
	ubiquitin ligase activity (By similarity). Regulator of Notch signaling, a signaling pathway	
	involved in cell-cell communications that regulates a broad spectrum of cell-fate	
	determinations. Mainly acts as a positive regulator of Notch, but it also acts as a negative	
	regulator, depending on the developmental and cell context. Mediates the antineural activity of	
	Notch, possibly by inhibiting the transcriptional activation mediated by MATCH1. Involved in	
	neurogenesis, lymphogenesis and myogenesis, and may also be involved in MZB (Marginal	
	zone B) cell differentiation. Promotes B-cell development at the expense of T-cell development,	
	suggesting that it can antagonize NOTCH1.	
Gene ID:	1840	
UniProt:	Q86Y01	
Pathways:	Notch Signaling	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
	-20 °C	
Storage:	-20 °C	

\vdash	land	linc
	iaria	11116

Expiry Date:

12 months