# antibodies -online.com





## anti-Deltex Homolog 1 antibody (AA 351-450) (Alexa Fluor 350)



Go to Product page

( )	11/0	K\ /	iew
	$\cup$	'I V/I	$I \cap VV$

Quantity:	100 μL
Target:	Deltex Homolog 1 (DTX1)
Binding Specificity:	AA 351-450
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Deltex Homolog 1 antibody is conjugated to Alexa Fluor 350
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human Deltex-1
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Cow
Purification:	Purified by Protein A.

#### **Target Details**

Target:	Deltex Homolog 1 (DTX1)
Alternative Name:	Deltex 1 (DTX1 Products)

### **Target Details**

9	
Background:	Synonyms: hDx-1, E3 ubiquitin-protein ligase DTX1, Protein deltex-1, Deltex1, hDTX1, DTX1
	Background: Functions as a ubiquitin ligase protein in vivo, mediating ubiquitination and
	promoting degradation of MEKK1, suggesting that it may regulate the Notch pathway via some
	ubiquitin ligase activity (By similarity). Regulator of Notch signaling, a signaling pathway
	involved in cell-cell communications that regulates a broad spectrum of cell-fate
	determinations. Mainly acts as a positive regulator of Notch, but it also acts as a negative
	regulator, depending on the developmental and cell context. Mediates the antineural activity of
	Notch, possibly by inhibiting the transcriptional activation mediated by MATCH1. Involved in
	neurogenesis, lymphogenesis and myogenesis, and may also be involved in MZB (Marginal
	zone B) cell differentiation. Promotes B-cell development at the expense of T-cell development,
	suggesting that it can antagonize NOTCH1.
Gene ID:	1840
UniProt:	Q86Y01
Pathways:	Notch Signaling
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C

			- 1	ı٠		
Н	la	n	М	П	n	
	а	11	u	П	1 1	Ч

Expiry Date:

12 months