antibodies -online.com





anti-DGAT1 antibody (AA 251-350) (AbBy Fluor® 555)



()	1 /	\sim	KI /	110	Νę
	1//	\vdash	I \/	1 ←	٠// ٢

Quantity:	100 μL	
Target:	DGAT1	
Binding Specificity:	AA 251-350	
Reactivity:	Human, Mouse, Rat, Dog	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This DGAT1 antibody is conjugated to AbBy Fluor® 555	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human DGAT1	
Isotype:	IgG	
Cross-Reactivity:	Dog, Human, Mouse, Rat	
Predicted Reactivity:	Pig,Horse	
Purification:	Purified by Protein A.	

Target Details

Target:	DGAT1	
Alternative Name:	ame: Dgat1/Diglyceride Acyltransferase (DGAT1 Products)	

Target Details

Background:	Synonyms: ARAT, DGAT, ARGP1, DIAR7, Diacylglycerol O-acyltransferase 1, ACAT-related gene	
	product 1, Acyl-CoA retinol O-fatty-acyltransferase, Retinol O-fatty-acyltransferase, Diglyceride	
	acyltransferase, DGAT1, AGRP1	
	Background: Catalyzes the terminal and only committed step in triacylglycerol synthesis by	
	using diacylglycerol and fatty acyl CoA as substrates. In contrast to DGAT2 it is not essential	
	for survival. May be involved in VLDL (very low density lipoprotein) assembly. In liver, plays a	
	role in esterifying exogenous fatty acids to glycerol. Functions as the major acyl-CoA retinol	
	acyltransferase (ARAT) in the skin, where it acts to maintain retinoid homeostasis and prevent	
	retinoid toxicity leading to skin and hair disorders.	
Gene ID:	8694	
UniProt:	075907	
Pathways:	Hormone Transport	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	