antibodies -online.com





anti-DOCK1 antibody (AA 465-550) (Alexa Fluor 555)



Go to Product page

\sim			
	N/P	r\/	i⊢₩

Quantity:	100 μL	
Target:	DOCK1	
Binding Specificity:	AA 465-550	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This DOCK1 antibody is conjugated to Alexa Fluor 555	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human DOCK1	
Isotype:	IgG	
Cross-Reactivity:	Mouse	
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Horse,Chicken	
Purification:	Purified by Protein A.	

Target Details

Target:	DOCK1
Alternative Name:	DOCK1 (DOCK1 Products)

Target Details

Synonyms: 180 kDa protein downstream of CRK, Ced5, Dedicator of Cytokinesis 1, Dedicator of cytokinesis protein 1, DOCK 1, DOCK-1, DOCK 180, DOCK1, DOCK1_HUMAN, DOCK180. Background: Involved in cytoskeletal rearrangements required for phagocytosis of apoptotic cells and cell motility. Functions as a guanine nucleotide exchange factor (GEF), which activates Rac Rho small GTPases by exchanging bound GDP for free GTP. Its GEF activity may be enhanced by ELMO1. Tissue specificity: Highly expressed in placenta, lung, kidney, pancreas	
and ovary. Expressed at intermediate level in thymus, testes and colon. 1793	
Signaling of Hepatocyte Growth Factor Receptor	
IF(IHC-P) 1:50-200	
IF(IHC-F) 1:50-200	
IF(ICC) 1:50-200	
For Research Use only	
Liquid	
1 μg/μL	
Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
ProClin	
This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
-20 °C	
Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
•	