antibodies -online.com





anti-DUSP7 antibody (AA 321-419) (Biotin)



Go to Product page

()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μL	
Target:	DUSP7	
Binding Specificity:	AA 321-419	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This DUSP7 antibody is conjugated to Biotin	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human DUSP7	
Isotype:	IgG	
Cross-Reactivity:	Human	
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Sheep,Pig,Horse,Chicken,Rabbit	
Purification:	Purified by Protein A.	

Target Details

Target:	DUSP7
Alternative Name:	DUSP7 (DUSP7 Products)

Target Details

Background:	Synonyms: Dual specicity phosphatase 7, Dual specicity protein phosphatase 7, Dual specicity			
	protein phosphatase PYST2, DUS7_HUMAN, Dusp7, Jnk Stimulatory Phosphatase 1, JSP1, MKPX, PYST2.			
	Background: Dual-specificity phosphatases (DUSPs) constitute a large heterogeneous			
	subgroup of the type I cysteine-based protein-tyrosine phosphatase superfamily. DUSPs are			
	characterized by their ability to dephosphorylate both tyrosine and serine/threonine residues.			
	DUSP7 belongs to a class of DUSPs, designated MKPs, that dephosphorylate MAPK (mitogen-			
	activated protein kinase) proteins ERK (see MIM 601795), JNK (see MIM 601158), and p38 (see			
	MIM 600289) with specificity distinct from that of individual MKP proteins. MKPs contain a			
	highly conserved C-terminal catalytic domain and an N-terminal Cdc25 (see MIM 116947)-like			
	(CH2) domain. MAPK activation cascades mediate various physiologic processes, including			
	cellular proliferation, apoptosis, differentiation, and stress responses			
Gene ID:	1849			
Pathways:	Neurotrophin Signaling Pathway, Activation of Innate immune Response, Toll-Like Receptors			
	Cascades			
Application Details				
Application Notes:	WB 1:300-5000			
	IHC-P 1:200-400			
	IHC-F 1:100-500			
Restrictions:	For Research Use only			
Handling				
Format:	Liquid			
Concentration:	1 μg/μL			
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.			
Preservative:	ProClin			
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be			
	handled by trained staff only.			
Storage:	-20 °C			
Storage Comment:	Store at -20°C for 12 months.			

\vdash	land	lına
	iaria	шц

Expiry Date:

12 months